


Cattle, sheep,
goats and pigs

Forticlina® Retard

動物藥入字第 5705 號



好適靈®

Oxytetracycline
廣效性抗生素

96 小時超長作用時間
抑制細菌蛋白質合成

INDICATIONS:


Cattle, sheep and goats:

- ▶ Pneumonia 肺炎
- ▶ Transport fever 運輸熱
- ▶ Pododermatitis and footrot 足皮炎
(complemented with local therapy)
- ▶ Diphtheria in calves 小牛的白喉
- ▶ Colibacillosis 大腸菌病
- ▶ Actinobacillosis 放射桿菌病
- ▶ Leptospirosis 細螺旋體病
- ▶ Anaplasmosis 紅血球胞子蟲病, 邊蟲病
- ▶ Metritis 子宮炎
- ▶ Mastitis 乳房炎

Pigs

- ▶ Colibacillosis 大腸菌病
- ▶ Pneumonia 肺炎
- ▶ Leptospirosis 細螺旋體病
- ▶ Metritis-mastitis-agalactia syndrome
(MMA) 子宮炎 - 乳房炎 - 無乳症
- ▶ Swine Erysipelas 豬丹毒





(成份) Oxytetracycline (dihydrate).....200mg
賦形劑.....加至.....1ml

(對象) 豬、牛、羊發生之對 Oxytetracycline 敏感之細菌性疾病

(用法用量) 依下列劑量行肌肉深部注射：

- 體重每 10 公斤使用 1ml
- 超過 100 公斤之動物應將劑量分半注射於兩個不同部位

(停藥期) 肉用 22 天 · 乳用 7 天

(包裝) 100ml × 30 瓶/盒，2 盒/箱

Forticlina® Retard



台灣總代理



佳力德實業有限公司
Jaritec Industrial Co.,Ltd.

http://www.jaritec.com.tw 地址：台北市信義路四段341號6樓之2

電話：02-2700-3163(代表號)

傳真：02-2325-7047

訂貨專線：0800-212-457

好畜蜜®



Committed to Animal Health & Nutrition